

The Post Rock Connection
Post Rock Extension District #1 Family and Consumer Sciences Column
Month of September 2024
By Jamie Rathbun, Family and Community Wellness Agent

Stay Safe, Eat Smart: Essential Tips to Prevent Foodborne Illness

When it comes to food safety, even a small mistake can lead to big consequences. With nearly 48 million Americans falling ill from foodborne illnesses each year, knowing how to handle food properly is more important than ever. Whether you're cooking at home or simply storing groceries, following a few simple steps can keep you and your loved ones safe from harmful bacteria. Let's break down the 4 essential tips to help you avoid becoming a statistic—and ensure that your meals are as safe as they are delicious.

Know the Symptoms

Consuming dangerous foodborne bacteria will usually cause illness within 1 to 3 days of eating the contaminated food. However, sickness can also occur within 20 minutes or up to 6 weeks later. Symptoms of foodborne illness can include: vomiting, diarrhea, and abdominal pain – and flu-like symptoms, such as fever, headache, and body ache.

Handle Foods Safely

Although most healthy people will recover from a foodborne illness within a short period of time, some can develop chronic, severe, or even life-threatening health problems. In addition, some people are at a higher risk for developing foodborne illness, including pregnant women, young children, older adults, and people with weakened immune systems (such as transplant patients and individuals with HIV/AIDS, cancer, or diabetes). To keep your family safer from food poisoning, **follow these four simple steps: clean, separate, cook, and chill.**

CLEAN: Wash hands and surfaces often

- Wash your hands with warm water and soap for at least 20 seconds before and after handling food and after using the bathroom, changing diapers, and handling pets.
- Wash your cutting boards, dishes, utensils, and counter tops with hot soapy water after preparing each food item.
- Consider using paper towels to clean up kitchen surfaces. If you use cloth towels, launder them often in the hot cycle.
- Rinse fresh fruits and vegetables under running tap water, including those with skins and rinds that are not eaten. Scrub firm produce with a clean produce brush.
- With canned goods, remember to clean lids before opening.

SEPARATE raw meats from other foods

- Separate raw meat, poultry, seafood, and eggs from other foods in your grocery shopping cart, grocery bags, and refrigerator.
- Use one cutting board for fresh produce and a separate one for raw meat, poultry, and seafood.
- Never place cooked food on a plate that previously held raw meat, poultry, seafood, or eggs unless the plate has been washed in hot, soapy water.
- Don't reuse marinades used on raw foods unless you bring them to a boil first.


COOK to the right temperature

- Color and texture are unreliable indicators of safety. Using a food thermometer is the only way to ensure the safety of meat, poultry, seafood, and egg products for all cooking methods. These foods must be cooked to a safe minimum internal temperature to destroy any harmful bacteria.
- Cook eggs until the yolk and white are firm. Only use recipes in which eggs are cooked or heated thoroughly.
- When cooking in a microwave oven, cover food, stir, and rotate for even cooking. If there is no turntable, rotate the dish by hand once or twice during cooking. Always allow standing time, which completes the cooking, before checking the internal temperature with a food thermometer.
- Bring sauces, soups and gravy to a boil when reheating.

CHILL: Refrigerate foods promptly

- Use an appliance thermometer to be sure the temperature is consistently 40° F or below and the freezer temperature is 0° F or below.
- Refrigerate or freeze meat, poultry, eggs, seafood, and other perishables within 2 hours of cooking or purchasing. Refrigerate within 1 hour if the temperature outside is above 90° F.
- Never thaw food at room temperature, such as on the counter top. There are three safe ways to defrost food: in the refrigerator, in cold water, and in the microwave. Food thawed in cold water or in the microwave should be cooked immediately.
- Always marinate food in the refrigerator.
- Divide large amounts of leftovers into shallow containers for quicker cooling in the refrigerator.

The Post Rock District of K-State Research and Extension serves Jewell, Lincoln, Mitchell, Osborne, and Smith Counties. Contact Jamie Rathbun at jrathbun@ksu.edu or by calling 785-524-4432. Stay connected with “Post Rock Extension” on Facebook, Twitter, Instagram, and YouTube. Our website is www.postrock.ksu.edu.



Post Rock Extension District #1

Beloit Office 115 S. Hersey Beloit, KS 67420 (785) 738-3597
Lincoln Office 108 W. Lincoln P.O. Box 8 Lincoln, KS 67455 (785) 524-4432
Mankato Office 307 N. Commercial Mankato, KS 66956 (785) 378-3174
Osborne Office 113 N. 1st Street Osborne, KS 67473 (785) 346-2521
Smith Center Office 218 S. Grant P.O. Box 287 Smith Center, KS 66967 (785) 282-6823

*Knowledge
for Life*

Kansas State University is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision, or hearing disability, contact a Post Rock District Office.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service
K-State Research and Extension is an equal opportunity provider and employer.